

# EFFECT OF SPATIAL DIFFERENTIATION OF PLANT COMMUNITIES ON PM<sub>2.5</sub> AND O<sub>3</sub> IN URBAN GREEN SPACES IN BEIJING, CHINA

Jianbin PAN<sup>✉</sup>, Shuyu CHEN, Nuo XU, Meijing CHENG, Xian WANG, Jingwen LAN, Rui WANG, Yajie WANG

*Department of Landscape Architecture, Beijing University of Civil Engineering and Architecture, Beijing, PR China*

## Highlights:

- the PM<sub>2.5</sub> concentrations are low in the areas of evergreen coniferous trees (ECP) and/or deciduous broadleaved trees (DBP) are the dominant species of tree-shrub-grass (TSG) and tree-shrub (TS);
- the O<sub>3</sub> concentration of all the plant community areas reaching the level of "low pollution";
- the AQI with PM<sub>2.5</sub>-O<sub>3</sub> value of compound concentration as the main parameter reaches the level of "moderate pollution", and the result that deserves further attention.

## Article History:

- received 30 August 2023
- accepted 26 August 2024

**Abstract.** Urban green space can improve the air quality of urban human settlements. This study aimed to investigate the spatial differences of air quality among the different plant community structures and types of urban park green spaces. We select 17 sample sites in Beijing Olympic Forest Park, and they are located in different areas of plant community structures and types. The study entailed an analysis of the interrelationships between the plant community structures, types, and PM<sub>2.5</sub>, O<sub>3</sub>, and PM<sub>2.5</sub>-O<sub>3</sub> compound data. The results showed that PM<sub>2.5</sub> was lower in tree-shrub-grass, tree-shrub, and tree-grass than in shrub-grass and grass plant community areas; PM<sub>2.5</sub> was lower in evergreen coniferous, mixed coniferous and broadleaved, and deciduous broadleaved plant communities than that in grass or shrub ones. In different plant community structures, types areas, O<sub>3</sub> was higher than 100 µg·m<sup>-3</sup>, and there were no significant differences among the plant community areas. The air quality index with PM<sub>2.5</sub>-O<sub>3</sub> composite pollution value as the main parameter reached the level of "moderate pollution", and the result that deserves further attention. The research results provide a basic scientific basis for the planning, design, and updating optimization of functional urban green spaces based on evidence-based design.

**Keywords:** air pollution, landscape architecture, urban green space, PM<sub>2.5</sub>-O<sub>3</sub>, spatial differentiation, evidence-based design.

<sup>✉</sup>Corresponding author. E-mail: [panjianbin@bucea.edu.cn](mailto:panjianbin@bucea.edu.cn)

## 1. Introduction

As urbanization in China increases and the population density in built-up areas continues to rise, environmental problems such as pollution aggravation have emerged in urban agglomerations such as Beijing-Tianjin-Hebei, Yangtze River Delta, Pearl River Delta, Chengdu-Chongqing, and especially in high-density urban areas such as the Beijing and Tianjin metropolitan region (Chen, 2020; Qi et al., 2017; Wu et al., 2021; Zhao & Xu, 2021; Feng et al., 2021; Liu et al., 2021; Liu et al., 2018; Xiao et al., 2022). In the last decade, major urban areas across China have experienced frequent exceedances of pollutant concentrations, which have seriously affected the health of residents, the quality of the urban microenvironment, and sustainable development (Fan et al., 2021; Li et al., 2022; Liu et al., 2021; Zhao et al., 2022). Fine particulate matter represented by

PM<sub>2.5</sub> and gaseous pollutants represented by O<sub>3</sub> have exceeded the standard concentration. Particularly in recent years, the Beijing-Tianjin-Hebei metropolitan region and other city clusters have shown the characteristics of multiple contamination of PM<sub>2.5</sub>, O<sub>3</sub>, and PM<sub>2.5</sub>-O<sub>3</sub> during the autumn-winter seasons (Cai et al., 2022; Chen, 2020; Li et al., 2022; Qin et al., 2019; Qu et al., 2018; Sheng et al., 2019; Wang et al., 2019; Zhao et al., 2018; Feng et al., 2021; Xiao et al., 2022). PM<sub>2.5</sub> mainly originates from industrial processes such as combustion and transport operations. It occurs mainly in the autumn-winter seasons and usually with significant spatial aggregation and diffusivity (Gao et al., 2020; Wang et al., 2019; Liu et al., 2018; Zhao et al., 2022). O<sub>3</sub> pollution mainly occurs in the summer and autumn. It has a certain homology and correlation with PM<sub>2.5</sub> pollution, because they share common precursors, nitrogen oxides (NO<sub>x</sub>) and volatile organic compounds (VOCs),

which produce  $O_3$  through photochemical reactions (Cai et al., 2022; Fishman & Crutzen, 1978). The spatio-temporal distribution features of  $NO_x$  and VOCs are different, because the mechanisms are not identical. Among the reduction (dissipation) and regional transmission (spill over) factors for both  $PM_{2.5}$  and  $O_3$ , larger-scale air movement (wind) and vegetation coverage are considered to be the most effective drivers (more than chemical energy use and population density) (Chen, 2020; Douglas et al., 2019; Qi et al., 2017; Liu et al., 2018; Wang et al., 2021). Previous studies have selected factors from the urban area scale, the street/block scale, and the landscape site scale to explain the underlying mechanisms for dissipating or reducing  $PM_{2.5}$ – $O_3$  compound pollution from the perspective of urban blue-green spatial system planning (Cai et al., 2022; Fan et al., 2021; Wu et al., 2021; Xiang et al., 2020), block/street morphology (He et al., 2023; Dai et al., 2020; Wang et al., 2021), and urban green space landscape spatial construction (Cai et al., 2022; Chen et al., 2019; King et al., 2014; Xing & Sun, 2022; Yin et al., 2022; Zhang et al., 2017). Some studies focusing on the plant community or individual scale of the plant landscape have concluded that  $PM_{2.5}$  concentration has a significant negative correlation with the amount of tri-dimensional green biomass of local plants (Zhu et al., 2019; Yan & Hong, 2019).

The study of  $PM_{2.5}$ – $O_3$  compound pollution at the national, regional, and city scales has mainly used measured data and spatial interpolation methods to describe the heterogeneity of  $PM_{2.5}$ – $O_3$  spatial distribution from a quantitative perspective, thus contributing to the formulation of joint prevention and control policies for cities and urban physical space planning. Studies at the landscape and urban green space scales have focused on the factors affecting  $PM_{2.5}$ – $O_3$  concentrations, thus providing a scientific basis for incremental urban spatial planning and urban regeneration based on functional green space design that reduces the intensity and scope of  $PM_{2.5}$ – $O_3$  pollution. However, in-depth quantitative studies on the spatial differentiation characteristics of  $PM_{2.5}$ – $O_3$  within larger urban green spaces are still relatively limited.

We continuously and dynamically monitored the microenvironment effects (such as reduction of airborne fungi and bacteria, cooling and humidification, and negative air ions) of the Beijing Olympic Forest Park from 2005 to 2022. Moreover, we have been monitoring  $PM_{2.5}$ – $O_3$  concentrations at regular intervals since 2013.

## 2. Study area and research methodology

### 2.1. Study area

The Beijing Olympic Forest Park (BOFP) is located in the Chaoyang District of Beijing, China, and covers a total area of 680  $hm^2$  (Figure 1); 450  $hm^2$  of the park is covered by more than 200 species of indigenous plants, forming a human-made near-natural forest system. Due to its large area and lush vegetation, the park plays a vital role in improving the urban microenvironment.

## 2.2. Research methods

### 2.2.1. Sample site setup

The chessboard sampling method was used to select 17 measured sample sites in BOFP (Figure 1). And the locations of the sample sites were carefully selected to ensure that they represent the air quality of typical environments (the sample sites of G, H and I are close to the urban road), and the others away from large areas of human activity (urban roads, squares, etc.) and to ensure representative vegetation types. 2–3 sample sites were tested for plant types, community structure, type, and typical landscape environment (Table 1).

Two comparison sample sites were selected, one 1 km from the south gate of BOFP (near the underground commercial plaza of the Olympic Park) and the other at the north side square of the North Fourth Ring Road (near the “Bird’s Nest” National Stadium, where there is a large amount of pavement, less green space, and activities involving dense crowds). Two comparison sample sites were not within the field of view of Figure 1.



**Figure 1.** Location of BOFP and 17 experimental sample sites in it

### 2.2.2. Test method

The test instrument comprised six outdoor air-quality testers (SWEMA TF-9, Sweden), which, in addition to  $PM_{2.5}$  and  $O_3$  measurements, could simultaneously collect and record  $PM_{10}$  and  $CO_2$  concentrations, air temperature, and relative humidity data (see Table 2 for information on instrument parameters). The experiment was conducted over 3 days during the period 10–25 August 2022, under a clear sky (no more than 30% cloud cover), with calm wind (within 3–4  $m\cdot s^{-1}$ ), while avoiding rainfall (in the event of rainfall, the experiment was postponed for 3 days). The biological characteristics of the plant community in the sample site area, such as Plant Height, Diameter at Breast Height, Crown Width, Forest Canopy, and Canopy Density of the Dominant Species in the quadrat were obtained by field research in the initial phase. A CI-110 Plant Canopy Im-

**Table 1.** Biological characteristics of sample sites in BOFP

Sample site	Plant community structure	Plant community type	Typical environment	Dominant species	Plant Height/m	Diameter at Breast Height/cm	Crown Width/m	Forest Canopy/m	Canopy Density/%
CK 1	–	–	–	–	–	–	–	–	–
CK 2	–	–	–	–	–	–	–	–	–
A	TG	DBP	DPC	<i>Populus tomentosa</i>	10~12	25~30	2.0~2.5	5.0~6.0	65
B	–	–	SPC	<i>Salix matsudana cv.pendula</i>	4.5~5.5	20~25	4.0~4.5	2.0~2.5	25
C	TSG	DBP	MPC	<i>Salix matsudana cv.pendula</i>	5.5~6.0	20~25	3.5~4.0	3.0~3.5	75
D	TSG	CBP	MPC	<i>Sabina chinensis, Sophora japonica</i>	3.5~4.0/ 5.5~6.0	20~25	2.0~2.5/ 4.5~5.0	1.5~2.0/ 2.5~3.0	85
E	T	CP	SPC	<i>Pinus tabulaeformis</i>	3.0~3.5	10~15	3.5~4.0	1.5~2.0	35
F	TG	DBP	DPC	<i>Salix matsudana</i>	7.0~8.0	20~25	4.5~5.0	3.0~4.0	85
G	SG	S	DPC	<i>Syringa oblata</i>	2.5~3.0	–	2.0~2.5	1.5~2.0	75
H	SG	S	DPC	<i>Caryopteris×clandonensis 'Worcester Gold'</i>	0.5~1.0	–	–	–	45
I	SG	S	DPC	<i>Euonymus japonicus</i>	0.5~1.0	–	–	–	45
J	SG	CP	DPC	<i>Pinus tabulaeformis</i>	4.5~5.0	10~15	2.5~3.0	2.0~2.5	55
K	SG	DBP	DPC	<i>Amygdalus triloba</i>	3.0~3.5	–	2.0~2.5	1.0~1.5	75
L	G	G	SPC	<i>Lawn and ground-cover</i>	–	–	–	–	75
M	TS	CP	DPC	<i>Pinus tabulaeformis</i>	3.5~4.0	10~15	2.5~3.0	1.5~2.0	95
N	TSG	CBP	MPC	<i>Populus tomentosa</i>	9.5~10.0	25~30	2.5~3.0	5.0~6.0	90
O	TSG	DBP	MPC	<i>Sophora japonica</i>	6.5~7.0	20~25	4.0~4.5	2.5~3.0	90
P	TG	DBP	DPC	<i>Ginkgo biloba</i>	4.5~5.0	15~25	2.5~3.0	2.0~2.5	75
Q	TG	DBP	DPC	<i>Sophora japonica, Fraxinus chinensis</i>	3.5~4.0	15~20	3.0~3.5/ 5~6	2~3/3~4	75

Note: (1) The sample site E is located in the "Heavenly Realm", which is 85 m above sea level, compared to other parkland sample sites at 43 m; (2) Plant community structures: tree-shrub-grass (TSG), tree-shrub (TS), tree-grass (TG), shrub-grass (SG), and grass/ground cover (G); plant community types: evergreen coniferous plant community (ECP), coniferous and broadleaved mixed plant community (CBP), deciduous broadleaved plant community (DBP), shrub (S), and grass/ground cover (G); typical environments: tree-shrub- grass multi-layer plant Community (MPC), tree-shrub, tree-grass, and shrub-grass double-layer plant community (DPC), grass/ground cover single-plant community (SPC), waterfront plant community (WPC), and waterfront square (WS); CK denotes the comparison sample.

age Analyzer was used to measure the leaf area index and other plant community quantification parameters.

**Table 2.** Test instrument parameters

Instrument	Items	Range of value	Accuracy
Outdoor air quality testers (SWEMA TF-9, Sweden)	PM <sub>2.5</sub>	0–1000 ug·m <sup>-3</sup>	±10% readout values
	O <sub>3</sub>	0–1200 ug·m <sup>-3</sup>	±20 ug·m <sup>-3</sup> +10% of readout values
	PM <sub>10</sub>	0–2000 ug·m <sup>-3</sup>	±10% readout values
	CO <sub>2</sub>	350–2000 PPM	±50 PPM+3% readout values
	Temperature	–20–50 °C	<±0.5 °C
	Relative Humidity	0–99%	<±3.5%

### 2.2.3. Data analysis methods

1) Statistical methods. The instrument automatically recorded and stored the measurement data from the sample sites. Measurements were taken from 8:00–18:00 (at 10 min intervals). Instantaneous value measurement, meanwhile, were taken in the morning, noon, and after-

noon at the following times: 8:50–9:10 (at 5 min intervals), 13:20–13:40, and 17:20–17:40. The arithmetic mean of the five times taken in each period was used as the value for morning, noon, and afternoon, respectively. The comparison sample data indicate the PM<sub>2.5</sub>–O<sub>3</sub> composite pollution concentration of the ambient background.

2) Air-quality evaluation method. According to the Ambient Air Quality Standards (GB 3095-2012, Chinese National Standards) (*Announcement on the Release of the Revision of Ambient Air Quality Standards (GB 3095-2012)*, 2018), released by the Ministry of Ecology and Environment in February 2012, the air-quality index (AQI) is a numerical value used to quantitatively describe the air quality: "excellent" (AQI ≤ 50), "good" (50 < AQI ≤ 100), "low pollution" (100 < AQI ≤ 150), "moderate pollution" (150 < AQI ≤ 200), "heavy pollution" (200 < AQI ≤ 300), and "severe pollution" (AQI > 300). This air-quality assessment is based on data recorded automatically at 10-min intervals for 10 consecutive hours from 8:00 to 18:00 (total of 76 recordings). The PM<sub>2.5</sub> and O<sub>3</sub> concentration values in this revised list can be used to evaluate the air-quality levels of the different plant communities in BOFP (Table 1). The formula for calculating the AQI is as follows:

$$AQI = \frac{AQI_{high} - AQI_{low}}{C_{high} - C_{low}} (C - C_{low}) + AQI_{low}, \quad (1)$$

where:  $C$  is the daily average of  $PM_{2.5}$  and  $O_3$  concentrations;  $I_{low}$  corresponds to the index limit of  $C_{low}$  (constant);  $I_{high}$  corresponds to the index limit of  $C_{high}$  (constant);  $C_{low}$  is less than or equal to the mass concentration limit of  $C$  (constant); and  $C_{high}$  is greater than or equal to the mass concentration limit of  $C$  (constant).

#### 2.2.4. Data processing

EXCEL 2019 was employed for organizing and conducting calculations about the average of  $PM_{2.5}$  and  $O_3$  concentrations, to facilitate subsequent comparisons. For significance analysis and one-way analysis of variance, SPSS 26 was utilized. Additionally, Origin 2023 was used to generate correlation bar charts and standard curves.

### 3. Results

#### 3.1. Spatial differentiation of $PM_{2.5}$ pollution

1)  $PM_{2.5}$  concentration and AQI within different plant community structure areas

Figure 2a shows the instantaneous  $PM_{2.5}$  value in the morning, noon, and afternoon within different plant community areas in BOFP. During the Morning period, the value of the grass/ground cover (G) plant community area was the highest (above  $50 \mu\text{g}\cdot\text{m}^{-3}$ ). The tree-shrub (TS) area was below  $50 \mu\text{g}\cdot\text{m}^{-3}$  (air quality "excellent"), with the lowest value. The  $PM_{2.5}$  concentrations in the tree-shrub-grass

(TSG) and the tree-shrub (TS) structure areas remained stable and decreased from noon to afternoon. However, the  $PM_{2.5}$  concentrations in the other plant community structure areas gradually increased, and all were higher than  $50 \mu\text{g}\cdot\text{m}^{-3}$ . Figure 2b compares the AQI values regarding  $PM_{2.5}$  concentrations in the different plant community structure areas. The  $PM_{2.5}$  concentrations in the areas with tree-shrub-grass (TSG), tree-grass (TG), and tree-shrub (TS) structures were below  $50 \mu\text{g}\cdot\text{m}^{-3}$  (air quality "excellent"), while the air quality in these areas with other community structures was just as high.

2)  $PM_{2.5}$  concentration and AQI within different plant community type areas

Figure 3a shows the instantaneous  $PM_{2.5}$  value in the morning, noon, and afternoon within different plant community type areas. The comparison between the three instantaneous values shows a slow increase in  $PM_{2.5}$  concentrations from morning to afternoon. The three instantaneous values of  $PM_{2.5}$  in the evergreen coniferous plant community (ECP), coniferous and deciduous broadleaf plant community (CBP, DBP) areas were all below  $50 \mu\text{g}\cdot\text{m}^{-3}$  (air quality "excellent"), while the instantaneous values at noon and afternoon in the shrub (S) area and the three instantaneous values in the grass/ground cover (G) area were above  $50 \mu\text{g}\cdot\text{m}^{-3}$ . Figure 3b compares AQI values within different plant community types areas. The  $PM_{2.5}$  concentrations in the evergreen coniferous (ECP), mixed coniferous and broadleaved (CBP), and deciduous broadleaf plant community (DBP) areas were below  $50 \mu\text{g}\cdot\text{m}^{-3}$ , and the air quality was assessed as "excellent".




Figure 2.  $PM_{2.5}$  instantaneous values ( $\mu\text{g}\cdot\text{m}^{-3}$ ) and AQI in different plant community structure areas




Figure 3.  $PM_{2.5}$  instantaneous values ( $\mu\text{g}\cdot\text{m}^{-3}$ ) and AQI in different plant community type areas

3)  $PM_{2.5}$  concentration and AQI within typical landscape environment areas

Figure 4a shows the instantaneous  $PM_{2.5}$  value in the morning, noon, and afternoon in the typical landscape environment of BOFP. The comparison of the three instantaneous values shows a slow increase from morning to afternoon in most areas, with only areas of the multi-layer plant community (MPC) showing an increase followed by a decrease. The  $PM_{2.5}$  concentrations in all landscape areas were below  $50 \mu\text{g}\cdot\text{m}^{-3}$  in the morning and below  $50 \mu\text{g}\cdot\text{m}^{-3}$  in the multi-layer (MPC) and waterfront plant community (WPC) areas (air quality "excellent"). In a combined comparison, the  $PM_{2.5}$  concentrations in BOFP were significantly lower than in the comparison sample sites. Figure 4b compares  $PM_{2.5}$  concentrations at the sample sites. The air quality of the single-layer (SPC) environment was rated as "good." In contrast, the other typical environments were rated as "excellent". On balance, the air quality indices in BOFP were better than those in the comparison samples.

### 3.2. Spatial differentiation of $O_3$ pollution

1)  $O_3$  concentration and AQI within different plant community structure areas

Figure 5a shows the  $O_3$  concentrations in the morning, noon, and afternoon in BOFP. The comparison between the three periods shows that the  $O_3$  concentrations in all plant community structure areas exhibited an increasing trend from morning to afternoon, with large increase from the morning to noon and a minor increase from noon to the afternoon.  $O_3$  concentrations in all areas of the plant community structure were below  $100 \mu\text{g}\cdot\text{m}^{-3}$  in the mornings




Figure 4.  $PM_{2.5}$  instantaneous values ( $\mu\text{g}\cdot\text{m}^{-3}$ ) and AQI in typical landscape environment areas

and above  $100 \mu\text{g}\cdot\text{m}^{-3}$  at noon and in the afternoons. Figure 5b shows a comparison of the  $O_3$  concentration in the different community structure areas. The difference in  $O_3$  concentration between the different community structures areas in BOFP is insignificant.  $O_3$  concentration is slightly lower than in the comparison sample sites, but not significantly different. All areas have "low Pollution."

2)  $O_3$  concentration and AQI within different plant community type areas

Figure 6a shows the  $O_3$  concentrations in the Morning, Noon, and Afternoon of the different plant community types areas in BOFP. The comparison between the three instantaneous values shows that  $O_3$  concentrations tended to increase from morning to afternoon, with higher initial values (above  $100 \mu\text{g}\cdot\text{m}^{-3}$ ) in the morning for evergreen coniferous (ECP), mixed coniferous and broadleaved (CBP) and deciduous broadleaf plant communities (DBP) and a slow increase from morning to afternoon; lower initial values (below  $100 \mu\text{g}\cdot\text{m}^{-3}$ ) in the morning for the two community types of shrub (S) and grass/ground Cover (G), and a large increase from morning to noon and a minor increase from noon to afternoon. Collectively, there were no significant differences between the Noon and Afternoon values of the  $O_3$  concentrations in all plant community types. Figure 6b shows a comparison of  $O_3$  concentrations in the different plant community types areas. Overall, no significant differences were found between the air quality of the different plant community types in BOFP. Although the air quality of BOFP is slightly higher than in the comparison sample sites (CK), there is still no significant difference, and the air quality is assessed as "low pollution."

3)  $O_3$  concentration and AQI within the typical landscape environment




Figure 5.  $O_3$  instantaneous values ( $\mu\text{g}\cdot\text{m}^{-3}$ ) and AQI in different plant community structure areas



Figure 7a shows the  $O_3$  concentrations in the morning, noon, and afternoon in the typical landscape environment area of BOFP. The comparison between the three instantaneous values shows that  $O_3$  concentrations in the typical landscape environment areas exhibited an increasing trend from morning to afternoon. However, the initial value concentrations were lower (below  $100 \mu\text{g}\cdot\text{m}^{-3}$ ) in the areas of the multi-layer (MPC), double-layer (DPC), and single-layer plant communities (SPC), with a large increase from the morning to noon and a minor increase from noon to afternoon. The noon and afternoon values of the  $O_3$  concentrations were above  $100 \mu\text{g}\cdot\text{m}^{-3}$  in all areas of the landscape environment. A comparison of the air quality assessment values, in terms of  $O_3$  concentrations, for the sample sites in typical landscape environments is shown in Figure 7b. There is no significant difference in air quality between the typical environmental areas. The air quality within BOFP is slightly higher than in the comparison sample sites, but not significantly different, all being “low pollution.”

### 3.3. Spatial differentiation of $\text{PM}_{2.5}\text{-O}_3$ compound pollution

Figure 8a to 8c compares the  $\text{PM}_{2.5}\text{-O}_3$  compound pollution AQI values of the structure and type of green space plant communities and typical landscape environments. The results show that the  $\text{PM}_{2.5}$  concentrations are low (below  $50 \mu\text{g}\cdot\text{m}^{-3}$ ) (air quality “excellent”) and the  $O_3$  concentrations are high (close to or above  $100 \mu\text{g}\cdot\text{m}^{-3}$ ) in each plant community structure, type, and typical landscape environment area. Thus, the  $\text{PM}_{2.5}\text{-O}_3$  concentration, as an air quality index, reached or exceeded  $150 \mu\text{g}\cdot\text{m}^{-3}$ , reach-

ing the level of “low pollution” or even “moderate pollution.” This result deserves further attention. In addition, the AQI ( $\text{PM}_{2.5}\text{-O}_3$  compound pollution) is relatively low in the areas of tree-shrub-grass (TSG), multi-layer (MPC), tree-shrub (TS), tree-grass (TG), double-layer (DPC), and waterfront plant communities (WPC).

## 4. Discussion

### 4.1. Spatially divergent correlations between green space plant communities and $\text{PM}_{2.5}$

In Figure 2a, 3a, and 4a, the slow increase in  $\text{PM}_{2.5}$  concentration during the three instantaneous periods of morning, noon, and afternoon. This increase in concentration may originate from the increase in dust and urban air pollutants caused by human activities, such as construction and transport operations, which are either direct or indirect sources of  $\text{PM}_{2.5}$  (e.g.,  $\text{NO}_x$  and VOCs, precursors of  $\text{PM}_{2.5}$ ). Changes in climatic factors in the urban area during the test time, such as a gradual increase in air temperature and decrease in relative humidity, also simultaneously lead to an increase in  $\text{PM}_{2.5}$  concentrations. This result further validates some scholars’ similar yet different findings, and so further research is needed (Cai et al., 2022; Xiao et al., 2022; Zhang et al., 2017; Zhao et al., 2014). In Figure 2a and 4a,  $\text{PM}_{2.5}$  concentrations in the area of the tree-shrub-grass plant community (TSG) increase and then decrease. This phenomenon is significant in the context of increasing  $\text{PM}_{2.5}$  concentrations. It is probably due to the more extensive leaf area index and tri-dimensional green biomass of the complex plant space of tree-shrub-grass




Figure 6.  $O_3$  instantaneous values ( $\mu\text{g}\cdot\text{m}^{-3}$ ) and AQI in different plant community type areas




Figure 7.  $O_3$  instantaneous values ( $\mu\text{g}\cdot\text{m}^{-3}$ ) and AQI in typical landscape environment areas

(TSG), which improves the urban microclimate conditions, for example, by increasing the horizontal and vertical air eddy and turbulence in the tree canopy, flushing fine particles into the canopy and further increasing the adhesion of fine particles to the plant leaf surface and facilitating plant. This may also be why the  $PM_{2.5}$  of urban green space was significantly lower than at the comparison sites (Cai et al., 2022; Fan et al., 2021; Niu et al., 2022; Sheng et al., 2019; Liu et al., 2021). Figure 2b, 3b and 4b show that the double-layered plant community (DPC) structure of tree-grass (TG), tree-shrub (TS) structure and evergreen coniferous (ECP), mixed coniferous and broadleaved (CBP) and deciduous broadleaf plant communities (DBP) as well as the waterfront plant community (WPC) areas can ensure turbulence between the forests, promoting horizontal and vertical air flow with canopy cover in the upper layers and enhancing the  $PM_{2.5}$  abatement effect of the tree canopy (Fan et al., 2021; Fishman & Crutzen, 1978; Jiang & Hong, 2021). The plants of evergreen coniferous (ECP) and mixed

coniferous plant community (CBP) in BOFP are still small, but the  $PM_{2.5}$  in these areas can be maintained within a certain concentration range, and the abatement effect of this community type deserves further attention (it is probably due to the dense branches and leaves, which are more conducive to promoting the dry deposition process of  $PM_{2.5}$ ). A layer of cover ensures turbulence in the forest, promoting horizontal and vertical airflow and further exerting the  $PM_{2.5}$  abatement effect of the tree canopy (Fan et al., 2021; Jiang & Hong, 2021).


#### 4.2. Spatially divergent correlations between green space plant communities and $O_3$

In Figure 5a, 6a, and 7a,  $O_3$  concentrations increased in the Morning, Noon, and Afternoon instantaneous for different plant community structures, types, and typical landscape environments. This increase was probably due to the gradual increase in solar radiation and the increased concentration of  $O_3$  precursor compounds ( $NO_x$  and VOCs), which favoured the production of higher  $O_3$  concentrations.  $O_3$  starting concentrations were low in all three periods. However, the increase in concentration was greater from morning to noon, while the increase tapered off from noon to afternoon. In the first period,  $O_3$  concentrations increased rapidly because the intensity of solar radiation increased rapidly to the highest value ( $O_3$ -producing conditions), and  $O_3$  precursor concentrations also reached a particularly high value ( $O_3$ -producing feedstock). In the second period,  $O_3$  concentrations increased to the highest value on the measurement day. Hence, the increase in concentration slowed. In Figure 5b, 6b, and 7b, there is no significant difference in air quality between the different structures, types of plant communities, and landscape environments (all reach the level of "low pollution"), probably because the green field plant communities themselves produce some amount of  $O_2$  through photosynthesis, but have no abatement on  $O_3$ . In Figure 7a, the morning  $O_3$  concentrations in the typical waterfront environment (WS, WPC) were higher than those in the typical plant community area and even higher than those in the comparison sample sites, probably because this environment is more favourable to  $O_3$  production due to the intense light in the morning (side shading only) (Fan et al., 2021). This result can be investigated in future studies (Cai et al., 2022; Qi et al., 2017).


#### 4.3. Spatially divergent correlations between green space plant communities and $PM_{2.5}$ - $O_3$ compound pollution

As shown in Figure 8a to 8c, the spatial variation of  $PM_{2.5}$  concentrations was significant for different plant community structures, types, and typical landscape environments in the urban green spaces. However,  $O_3$  concentrations did not show significant spatial variation and were high (all above  $100 \mu g \cdot m^{-3}$ ). Based on the results above, the air quality was assessed as "moderate pollution" because of the high contribution of  $O_3$  to the  $PM_{2.5}$ - $O_3$  compound pollution (over 70%). Most researchers now believe that  $PM_{2.5}$  pollution

a) Plant community structure areas



b) Plant community type areas



c) Typical landscape environment areas




Figure 8. AQI ( $PM_{2.5}$ - $O_3$  combined concentration)

and O<sub>3</sub> pollution are similar and homogenous (Fan et al., 2021; Gao et al., 2020). Air movement (wind) is considered the most effective mechanism for abatement both, followed by factors such as green vegetation. However, it can be inferred from the results of this study that green vegetation areas can abate PM<sub>2.5</sub> to varying degrees, but no significant abatement of O<sub>3</sub> was observed. Some scholars therefore believe that regional coordination, joint prevention, and control mechanisms should be adopted for PM<sub>2.5</sub>–O<sub>3</sub> compound pollution. The control and reduction of PM<sub>2.5</sub> and O<sub>3</sub> precursors (industrial processes and fossil energy-using urban traffic) and the creation of mechanisms for the abatement and rapid dispersion of PM<sub>2.5</sub>–O<sub>3</sub> pollution (improvement and optimization of the neighbourhood wind environment and the formation of “Urban Wind Corridors” in general planning) are central to this joint prevention and control strategy (Li et al., 2022; Dai et al., 2020; Feng et al., 2021; Wang et al., 2021; Xiao et al., 2022).

## 5. Conclusions

This study aimed to provide a scientific basis for the renewal and optimization of urban green spaces. The results allow us to draw the following conclusions:

- The PM<sub>2.5</sub> concentrations are low in the areas of evergreen coniferous trees (ECP) and/or deciduous broadleaved trees (DBP) are the dominant species of tree-shrub-grass (TSG) and tree-shrub (TS).
- The O<sub>3</sub> concentration of all the plant community areas reaching the level of “low pollution”.
- The AQI with PM<sub>2.5</sub>–O<sub>3</sub> value of compound concentration as the main parameter reaches the level of “moderate pollution”, and the result that deserves further attention.

The findings may not be universal, however, due to the study’s significant spatial and temporal heterogeneity. In particular, given the openness and complex system of urban green spaces, we modelled the spatial composition of green space plant communities but may have overlooked or ignored the heterogeneous information that has an impact on the results. Furthermore, three instantaneous concentration measurements from the morning, noon, and afternoon were used as the daily average concentrations, whereas the data underlying the air-quality evaluation were taken at 08:00. The data were automatically recorded at 10 min intervals for 10 consecutive hours from 08:00 to 18:00, and there may be some differences in accuracy between the two. The conclusions are therefore tentative and should be subject to more reviews and test.

## Funding

This work was supported by the National Natural Science Foundation of China Project, under the grant titled “Basic research for the parametric design of green landscape of outdoor environment based on negative-aeroion comfort” (51641801).

## Author contributions

Jianbin Pan: conceptualization, methodology, software, validation, formal analysis, resources, data curation, visualization, supervision, project administration, funding acquisition; Shuyu Chen: methodology, software, validation, investigation, resources, data curation, writing-original draft preparation, writing-review and editing, visualization, funding acquisition; Nuo Xu: validation, formal analysis, data curation, writing-original draft preparation, writing-review and editing; Meijing Cheng: investigation; Xian Wang: investigation; Jingwen Lan: investigation; Rui Wang: data curation; Yajie Wang: data curation. All authors have read and agreed to the published version of the manuscript.

## Disclosure statement

The authors declare no conflict of interest.

## References

- Cai, L., Zhuang, M., & Ren, Y. (2022). Spatiotemporal characteristics of NO<sub>2</sub>, PM<sub>2.5</sub> and O<sub>3</sub> in a coastal region of southeastern China and their removal by green spaces. *International Journal of Environmental Health Research*, 32(1), 1–17. <https://doi.org/10.1080/09603123.2020.1720620>
- Chen, A. S. Z. (2020, August 28–30). *Temporal distribution characteristics of PM<sub>2.5</sub> in Beijing and its influential factors* [Conference presentation]. IOP Conference Series: Earth and Environmental Science, 6th International Conference on Energy, Environment and Materials Science (EEMS), Hulun Buir, China. <https://doi.org/10.1088/1755-1315/585/1/012041>
- Chen, M., Dai, F., Yang, B., & Zhu, S. W. (2019). Effects of neighborhood green space on PM<sub>2.5</sub> mitigation: Evidence from five megacities in China. *Building and Environment*, 156, 33–45. <https://doi.org/10.1016/j.buildenv.2019.03.007>
- Dai, F., Chen, M., Wang, M., Zhu, S. W., & Fu, F. (2020). Effect of urban block form on reducing particulate matter: A case study of Wuhan. *Chinese Landscape Architecture*, 36(03), 109–114. <https://doi.org/10.19775/j.cla.2020.03.0109>
- Douglas, A. N. J., Irga, P. J., & Torpy, F. R. (2019). Determining broad scale associations between air pollutants and urban forestry: A novel multifaceted methodological approach. *Environmental Pollution*, 247, 474–481. <https://doi.org/10.1016/j.envpol.2018.12.099>
- Fan, S. X., Zhang, M. Y., Li, Y. L., Li, K., & Dong, L. (2021). Impacts of composition and canopy characteristics of plant communities on microclimate and airborne particles in Beijing, China. *Sustainability*, 13(9), Article 4791. <https://doi.org/10.3390/su13094791>
- Feng, N., Tang, M. X., Li, M. L., Chen, Y., Cao, L. M., He, L. Y., & Huang, X. F. (2021). Research on the influence of VOCs on the coupling generation of PM<sub>2.5</sub> and O<sub>3</sub> in Shenzhen. *China Environmental Science*, 41(01), 11–17. <https://doi.org/10.19674/j.cnki.issn1000-6923.2021.0002>
- Fishman, J., & Crutzen, P. J. (1978). The origin of ozone in the troposphere. *Nature*, 274, 855–858. <https://doi.org/10.1038/274855a0>
- Gao, T., Liu, F., Wang, Y., Mu, S., & Qiu, L. (2020). Reduction of atmospheric suspended particulate matter concentration and influencing factors of green space in urban forest park. *Forests*, 11(9), Article 950. <https://doi.org/10.3390/f11090950>



- He, H. Y., Zhu, Y. S., Liu, L., Du, J., Liu, L. R., & Liu, J. (2023). Effects of roadside trees three-dimensional morphology characteristics on traffic-related PM<sub>2.5</sub> distribution in hot-humid urban blocks. *Urban Climate*, 49, Article 101448. <https://doi.org/10.1016/j.uclim.2023.101448>
- Jiang, R. S., & Hong, B. (2021). Spatio-temporal distribution characteristics of PM<sub>2.5</sub> and PM<sub>10</sub> and residents' exposure risk assessment in residential outdoor open spaces. *Chinese Landscape Architecture*, 37(08), 121–126. <https://doi.org/10.19775/j.cla.2021.08.0121>
- King, K. L., Johnson, S., Kheirbek, I., Lu, J. W. T., & Matte, T. (2014). Differences in magnitude and spatial distribution of urban forest pollution deposition rates, air pollution emissions, and ambient neighborhood air quality in New York City. *Landscape and Urban Planning*, 128, 14–22. <https://doi.org/10.1016/j.landurbplan.2014.04.009>
- Li, Z. Y., Xie, M. M., Wang, H. H., Chen, B., Wu, R. R., & Chen, Y. (2022). The spatiotemporal heterogeneity of the relationship between PM<sub>2.5</sub> concentrations and the surface urban heat island effect in Beijing, China. *Progress in Physical Geography-Earth and Environment*, 46(1), 84–104. <https://doi.org/10.1177/03091333211033209>
- Liu, C., Jin, M. Y., Zhu, X. H., & Peng, Z. R. (2021). Review of patterns of spatiotemporal PM<sub>2.5</sub> driving factors, methods evolution and urban planning implications. *Journal of Human Settlements in West China*, 36(04), 9–18. <https://doi.org/10.13791/j.cnki.hsfwest.20210402>
- Liu, H., Fang, C., Huang, X., Zhu, X., Zhou, Y., Wang, Z., & Zhang, Q. (2018). The spatial-temporal characteristics and influencing factors of air pollution in Beijing-Tianjin-Hebei urban agglomeration. *Acta Geographica Sinica*, 73(1), 177–191. <https://doi.org/10.11821/dlxb201801015>
- Ministry of Ecology and Environment. (2018). *Announcement on the release of the revision of Ambient Air Quality Standards (GB 3095-2012)*. [http://www.mee.gov.cn/gkml/sthjbgw/sthjbgg/201808/t20180815\\_451398.htm](http://www.mee.gov.cn/gkml/sthjbgw/sthjbgg/201808/t20180815_451398.htm)
- Niu, X., Li, Y., Li, M. N., Zhang, T., Meng, H., Zhang, Z., Wang, B., & Zhang, W. K. (2022). Understanding vegetation structures in green spaces to regulate atmospheric particulate matter and negative air ions. *Atmospheric Pollution Research*, 13(9), Article 101534. <https://doi.org/10.1016/j.apr.2022.101534>
- Qi, B., Niu, Y., Du, R., Yu, Z., Ying, F., Xu, H., Hong, S., & Yang, H. (2017). Characteristics of surface ozone concentration in urban site of Hangzhou. *China Environmental Science*, 37(02), 443–451.
- Qin, H. Q., Hong, B., Jiang, R. S., Yan, S. S., & Zhou, Y. H. (2019). The effect of vegetation enhancement on particulate pollution reduction: CFD Simulations in an urban park. *Forests*, 10(5), Article 373. <https://doi.org/10.3390/f10050373>
- Qu, Y. W., Wang, T. J., Cai, Y. F., Wang, S. K., Chen, P. L., Li, S., Li, M. M., Yuan, C., Wang, J., & Xu, S. C. (2018). Influence of atmospheric particulate matter on ozone in Nanjing, China: Observational study and mechanistic analysis. *Advances in Atmospheric Sciences*, 35(11), 1381–1395. <https://doi.org/10.1007/s00376-018-8027-4>
- Sheng, Q. Q., Zhang, Y. L., Zhu, Z. L., Li, W. Z., Xu, J. Y., & Tan, R. (2019). An experimental study to quantify road greenbelts and their association with PM<sub>2.5</sub> concentration along city main roads in Nanjing, China. *Science of the Total Environment*, 667, 710–717. <https://doi.org/10.1016/j.scitotenv.2019.02.306>
- Wang, P., Guo, H., Hu, J., Kota, S. H., Ying, Q., & Zhang, H. (2019). Responses of PM<sub>2.5</sub> and O<sub>3</sub> concentrations to changes of meteorology and emissions in China. *Science of the Total Environment*, 662, 297–306. <https://doi.org/10.1016/j.scitotenv.2019.01.227>
- Wang, W., Cheng, X. Y., Hu, C., Xia, S. H., & Wang, T. (2021). Spatio-temporal distribution characteristics of PM<sub>2.5</sub> and air quality evaluation in urban street canyons: Take Changhuai Street in Hefei as an example. *Ecology and Environmental Sciences*, 30(11), 2157–2164. <https://doi.org/10.16258/j.cnki.1674-5906.2021.11.006>
- Wu, J., Wang, Y., Liang, J., & Yao, F. (2021). Exploring common factors influencing PM<sub>2.5</sub> and O<sub>3</sub> concentrations in the Pearl River Delta: Tradeoffs and synergies. *Environmental Pollution*, 285, Article 117138. <https://doi.org/10.1016/j.envpol.2021.117138>
- Xiang, S., Liu, J., Tao, W., Yi, K., Xu, J., Hu, X., Liu, H., Wang, Y., Zhang, Y., Yang, H., Hu, J., Wan, Y., Wang, X., Ma, J., Wang, X., & Tao, S. (2020). Control of both PM<sub>2.5</sub> and O<sub>3</sub> in Beijing-Tianjin-Hebei and the surrounding areas. *Atmospheric Environment*, 224, Article 117259. <https://doi.org/10.1016/j.atmosenv.2020.117259>
- Xiao, Z. M., Xu, H., Gao, J. Y., Cai, Z. Y., Bi, W. K., Li, P., Yang, N., Deng, X. W., & Ji, Y. F. (2022). Characteristics and sources of PM<sub>2.5</sub>-O<sub>3</sub> compound pollution in Tianjin. *Environmental Science*, 43(03), 1140–1150. <https://doi.org/10.13227/j.hjxk.202108164>
- Xing, Q. F., & Sun, M. P. (2022). Characteristics of PM<sub>2.5</sub> and PM<sub>10</sub> spatio-temporal distribution and influencing meteorological conditions in Beijing. *Atmosphere*, 13(7), Article 1120. <https://doi.org/10.3390/atmos13071120>
- Yan, S. S., & Hong, B. (2019). PM<sub>2.5</sub> concentration distribution characteristics in different landscape spaces and influencing factors in urban park. *Landscape Architecture*, 26(07), 101–106. <https://doi.org/10.14085/j.fjyl.2019.07.0101.06>
- Yin, Z., Zhang, Y. X., & Ma, K. M. (2022). Evaluation of PM<sub>2.5</sub> retention capacity and structural optimization of urban park green spaces in Beijing. *Forests*, 13(3), Article 415. <https://doi.org/10.3390/f13030415>
- Zhang, K., Meng, F., Li, X. Y., Zhou, J., & Cui, K. Q. (2017). Effect of landscape plants on the concentration of PM<sub>2.5</sub> from vehicle emission. *Ecology and Environmental Sciences*, 26(06), 1009–1016. <https://doi.org/10.16258/j.cnki.1674-5906.2017.06.014>
- Zhao, A. Z., Xiang, K. Z., Liu, X. F., & Zhang, X. R. (2022). Spatio-temporal evolution patterns of PM<sub>2.5</sub> and relationship with urban expansion in Beijing-Tianjin-Hebei urban agglomeration from 2000 to 2018. *Environmental Science*, 43(05), 2274–2283. <https://doi.org/10.13227/j.hjxk.202109226>
- Zhao, C. X., Wang, Y. Q., Wang, Y. J., Zhang, H. L., & Zhao, B. Q. (2014). Temporal and spatial distribution of PM<sub>2.5</sub> and PM<sub>10</sub> pollution status and the correlation of particulate matters and meteorological factors during winter and spring in Beijing. *Environmental Science*, 35(02), 418–427. <https://doi.org/10.13227/j.hjxk.2014.02.013>
- Zhao, H., Zheng, Y., & Li, C. (2018). Spatiotemporal distribution of PM<sub>2.5</sub> and O<sub>3</sub> and their interaction during the summer and winter seasons in Beijing, China. *Sustainability*, 10(12), Article 4519. <https://doi.org/10.3390/su10124519>
- Zhao, S., & Xu, Y. (2021). Exploring the dynamic spatio-temporal correlations between PM<sub>2.5</sub> emissions from different sources and urban expansion in Beijing-Tianjin-Hebei Region. *International Journal of Environmental Research and Public Health*, 18(2), Article 608. <https://doi.org/10.3390/ijerph18020608>
- Zhu, C. Y., Przybysz, A., Chen, Y. R., Guo, H. J., Chen, Y. Y., & Zeng, Y. Z. (2019). Effect of spatial heterogeneity of plant communities on air PM<sub>10</sub> and PM<sub>2.5</sub> in an urban forest park in Wuhan, China. *Urban Forestry & Urban Greening*, 46, Article 126487. <https://doi.org/10.1016/j.ufug.2019.126487>